

ENRAILIFE

Biostimulant with amino acids that favors the development of root systems in horticultural crops, ornamentals and fruit-trees.

GUARANTEED RICHNESS

Free amino acids: 8% = (9,2% m/v)	Molybdenum (Mo) soluble in water: 0,001%
Total nitrogen (N): 5,6%	Boron(B) soluble in water: 0,01%
Organic nitrogen (N): 1,6%	EDTA chelated zinc (Zn): 0,05%
Ureic nitrogen (N): 4%	Organic matter: 25% = (29% m/v)
EDTA chelated copper (Cu): 0,05%	EDTA chelated manganese (Mn): 0,05%

GENERAL FEATURES

ENRAILIFE is a **biostimulant** that contains natural amino acids, low molecular weight peptides, vegetal polysaccharides and chelated microelements. This product is specially recommended to be applied via fertigation. Its balanced composition assures the maximum availability of nutrients for the plant. When amino acids are grouped, the proteins begin to form. **ENRAILIFE** contains amino acids that, when absorbed, save the plant the task of synthesizing them. Those savings allow the plants to take full advantage of their genetic potential.

The amino acids contained in this product are absorbed rapidly by the root system and they are transported to the organs of more activity. Due to its high content in organic matter (25%), the regular application of **ENRAILIFE** improves the soil's physicochemical properties and increases the development of the root system.

Manufactured by:

AGROQUIMICOS Y NUTRIENTES, S.L.

RECOMMENDATIONS FOR USE [®]

Dose and instructions for use

ENRAILIFE is a fertilizer developed for its application via fertigation:

- Transplant of **horticultural plants, flowers and ornamental**: from 3 to 5 liters per hectare after the transplant, then another time using the same dose.
- **Fruit-trees** (citrus, olive-tree, grapevines, apple-tree, etc.): using from 3 to 4 liters per hectare and 4 applications throughout the vegetative cycle.

Application season

ENRAILIFE can be used in **all types of crops** during the rooting and the growth stages. Its application for **horticultural plants** is advisable 7-14 days after the transplant and another application after 15-20 days. In **citrus and fruit-trees**, it can be applied throughout the vegetative cycle in order to increase the development of the root system and increase the assimilation of nutrients.

Presentation

Drums of 5 and 20 liters and containers of 1000 liters.